COVID-19 Screening Classifications and Definitions for People Experiencing Homelessness

The following chart and definitions are for reference by homeless assistance and medical providers responsible for screening and triaging individuals who may be confirmed or suspected of having COVID-19. The information included here is not medical advice and is subject to change based on CDC, local public health, or other government guidance and directive.

Screening	Definition	Intervention
Classification		
COVID-19 Confirmed	Individual with positive COVID test within prior	9-1-1 and/or appropriate medical
or Symptomatic,	14 days or currently symptomatic and	facility
Severe Respiratory	experiencing severe related symptoms and	Providers use full PPE
Distress	medical distress.	Mask individual
COVID-19 Confirmed	Individual with positive COVID test within prior	Isolation at appropriate medical facility
or Symptomatic, <u>Not</u>	14 days (with or without symptoms), OR with	Providers use full PPE
Appropriate for SIQ	mild-to-moderate COVID symptoms within	Mask individual
options	past 24 hours not due to other underlying	• Monitor for more severe symptoms
	conditions, AND who has significant health	
	and/or ADL needs that cannot be met by	
	existing SIQ options.	
COVID-19 Confirmed	Individual with positive COVID test within prior	Isolation at SIQ facility, if available
or Symptomatic,	14 days (with or without symptoms) OR with	Providers use full PPE
Appropriate for SIQ	mild-to-moderate COVID symptoms within	Mask individual
options	past 24 hours not due other underlying	Monitor for more severe symptoms
	conditions, AND who <u>does not have</u> significant	
	health and/or ADL needs that cannot be met	
	by existing SIQ options.	
COVID-19 Exposed,	Individual without mild-to-moderate	Quarantine at appropriate medical
Presumed Positive,	respiratory symptoms in the last 72 hours who	facility
Not Appropriate for	had close contact with someone confirmed for	Providers use basic PPE (mask, face
SIQ options	COVID-19 in the past 14 days, AND who has	shield, gloves)
	significant health and/or ADL needs that cannot be met by existing SIQ options.	Mask individual
COV/ID 10 Frances d		Monitor for more severe symptoms
COVID-19 Exposed,	Individual without mild-to-moderate	Quarantine at SIQ facility, if available
Presumed Positive, <u>Appropriate</u> for SIQ	respiratory symptoms in the last 72 hours who had close contact with someone confirmed for	Providers use basic PPE (mask, face shield, glayes)
options	COVID-19 in the past 14 days, AND who does	shield, gloves)Mask individual
options	<u>not have</u> significant health and/or ADL needs	
	that cannot be met by existing SIQ options.	Monitor for more severe symptoms
COVID-19 Presumed	Individual not confirmed COVID-19 and	Quasi-Quarantine as much as possible
Negative, Low Risk	without mild-to-moderate respiratory	at existing shelter or current location.
and/or Vulnerability	symptoms in the last 72 hours; may have had	 Providers use basic PPE (mask, face
,	incidental, non-close contact with someone	shield, gloves)
	confirmed for COVID-19 in the past 14 days.	Mask individual
	Individual <u>does not</u> have high risk or	Monitor for more severe symptoms
	vulnerability of severe illness from COVID-19.	
COVID-19 Presumed	Individual not confirmed COVID-19 and	Quasi-Quarantine as much as possible
Negative, At-Risk or	without mild-to-moderate respiratory	at existing shelter or current location.
Vulnerable	symptoms in the last 72 hours; may have had	• Providers use basic PPE (mask, face
	incidental, non-close contact with someone	shield, gloves)
	confirmed for COVID-19 in the past 14 days.	Mask individual
	Individual <u>has</u> high risk or vulnerability of	• Monitor for more severe symptoms
	severe illness from COVID-19.	

Rev. 03-31-2020

COVID-19 Symptoms

- Mild-to-moderate symptoms:
 - Fever ≥100.4°F
 - New cough developed in last 14 days
 - o Shortness of breath
- Additional non-severe symptoms (not all inclusive):
 - Unexplained muscle aches
 - o Sore throat
 - o Runny nose
 - Nasal congestion
 - o Sneezing
- Severe respiratory distress symptoms requiring immediate medical attention (not all inclusive):
 - o Trouble breathing
 - Persistent pain or pressure in the chest
 - o New confusion or inability to arouse
 - o Bluish lips or face
 - o Extreme fatigue

Close Contact ("Exposure") includes:

- Living in the same household as a sick person with COVID-19
- Caring for a sick person with COVID-19
- Being within 6 feet of a sick person with COVID-19 for about 10 minutes
- Being in direct contact with secretions from a sick person with COVID-19 (e.g., being coughed on, kissing, sharing utensils, etc.)

High Risk/Vulnerability of Severe Illness from COVID-19 include individuals who:

- Are over 60 years of age
- Have underlying medical conditions
- Are pregnant

For questions regarding this, please contact Erin Maus, Community Shelter Board at <u>emaus@csb.org</u>.